

Course of **Analysis I**

Chapter V : LIMITED DEVELOPMENT

Academic Year 2025-2026

Contents

1	Limited developments	2
1.1	Notations	2
1.2	Definition and existence	2
1.3	The Limited development of Common functions at 0 up to Order n	4
2	Operations on limited developments	5
2.1	Sum and Product	5
2.2	Composition	6
2.3	Quotient	7
2.4	Integration	8
3	Applications of Limit Developments	9
3.1	Limit Calculation	9
3.2	Continuity and Differentiability from a limited development	9

In this chapter, for any given function, we will find the polynomial of degree n that best approximates the function around a specific point. More precisely, we aim to decompose any sufficiently regular function f around a given point a as follows:

$$f(x) = T_n(x) + R_n(x);$$

where T_n is a polynomial of degree n and R_n is a function that satisfies $\lim_{x \rightarrow a} R_n(x) = 0$.

Decomposing a function in this way around a point a is known as performing a limited development (LD) of the function at point a to order n . The polynomial T_n is called the polynomial part of the LD, and the function R_n is referred to as the remainder of the expansion.

1 Limited developments

1.1 Notations

Let $I \subset \mathbb{R}$ be an open interval, $f : I \rightarrow \mathbb{R}$ a function, and $n \in \mathbb{N}^*$. - If f is differentiable and if the function $f' : I \rightarrow \mathbb{R}$ is also differentiable, we denote $f'' = (f')'$ as the second derivative of f . More generally, for any $n \in \mathbb{N}$, we denote

$$f^{(0)} = f, f^{(1)} = f', f^{(2)} = f'', \dots, f^{(n+1)} = (f^{(n)})'$$

- f is of class C^0 on I if f is continuous on I . We write $f \in C^0(I)$.
- f is of class C^n on I if f is n -times differentiable on I and $f^{(n)}$ is continuous on I . We write $f \in C^n(I)$.
- f is of class C^∞ on I if $\forall n \in \mathbb{N}, f \in C^n(I)$. We write $f \in C^\infty(I)$. Note that for any $n \in \mathbb{N}$, $C^\infty(I) \subset C^n(I)$.
- For any $n \in \mathbb{N}^*$, the factorial of n is defined by $n! = 1 \times 2 \times \dots \times (n-1) \times n$ with the convention $0! = 1$.

1.2 Definition and existence

Let I be an open interval and $f : I \rightarrow \mathbb{R}$ a function.

Definition 1. For $a \in I$ and $n \in \mathbb{N}$, we say that f admits a limited development at point a to order n if there exist $n+1$ real constants c_0, c_1, \dots, c_n and a function $\varepsilon : I \rightarrow \mathbb{R}$ such that $\lim_{x \rightarrow a} \varepsilon(x) = 0$ and, for every x in I ,

$$f(x) = \underbrace{c_0 + c_1(x-a) + c_2(x-a)^2 + \dots + c_n(x-a)^n}_{\text{Polynomial part of the LD}} + \underbrace{(x-a)^n \varepsilon(x)}_{\text{Remainder of the LD}}.$$

Proposition 1. Let I be an open interval and $a \in I$. If f is of class C^n on I , then f admits a limited development of order n at the point a given by the following formula known as the Taylor-Young formula:

$$f(x) = f(a) + f'(a)(x-a) + \frac{f''(a)}{2!}(x-a)^2 + \dots + \frac{f^{(n)}(a)}{n!}(x-a)^n + (x-a)^n \varepsilon(x),$$

where $\lim_{x \rightarrow a} \varepsilon(x) = 0$.

Remark 1. The limited development of a function f of order n at the point 0 is written as follows using the Taylor-Young formula:

$$f(x) = f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \dots + \frac{f^{(n)}(0)}{n!}x^n + x^n\epsilon(x)$$

where $\lim_{x \rightarrow 0} \epsilon(x) = 0$.

Example 1. Let $a \in \mathbb{R}$ and $n \in \mathbb{N}^*$. Give the limited development at the point a to order n of the function $f(x) = e^x$.

$$f(x) = e^x, \quad f'(x) = e^x, \quad f''(x) = e^x, \quad \dots, \quad f^{(n)}(x) = e^x.$$

Evaluating each at $x = a$, we obtain:

$$f(a) = e^a, \quad f'(a) = e^a, \quad f''(a) = e^a, \quad \dots, \quad f^{(n)}(a) = e^a.$$

Thus, the limited development of e^x at $x = a$ to order n is:

$$\begin{aligned} e^x &= f(a) + f'(a)(x-a) + \frac{f''(a)}{2!}(x-a)^2 + \dots + \frac{f^{(n)}(a)}{n!}(x-a)^n + (x-a)^n\epsilon(x). \\ &= e^a + e^a(x-a) + \frac{e^a}{2!}(x-a)^2 + \dots + \frac{e^a}{n!}(x-a)^n + (x-a)^n\epsilon(x). \\ &= e^a \left(1 + (x-a) + \frac{(x-a)^2}{2!} + \dots + \frac{(x-a)^n}{n!} \right) + (x-a)^n\epsilon(x), \end{aligned}$$

where $\lim_{x \rightarrow 0} \epsilon(x) = 0$.

Example 2. Give the limited development of $f(x) = \sin x$ of order 5 at the point 0 .

$$f(x) = \sin x, \quad f'(x) = \cos x, \quad f''(x) = -\sin x, \quad f^{(3)}(x) = -\cos x, \quad f^{(4)}(x) = \sin x, \quad f^{(5)}(x) = \cos x$$

Evaluating each at $x = a$, we obtain:

$$f(0) = 0, \quad f'(0) = 1, \quad f''(0) = 0, \quad f^{(3)}(0) = -1, \quad f^{(4)}(0) = 0, \quad f^{(5)}(0) = 1.$$

Then

$$\begin{aligned} \sin(x) &= f(0) + \frac{f'(0)}{1!}x + \frac{f''(0)}{2!}x^2 + \frac{f^{(3)}(0)}{3!}x^3 + \frac{f^{(4)}(0)}{4!}x^4 + \frac{f^{(5)}(0)}{5!}x^5 + x^5\epsilon(x), \\ &= x - \frac{x^3}{3!} + \frac{x^5}{5!} + x^5\epsilon(x). \end{aligned}$$

where $\lim_{x \rightarrow 0} \epsilon(x) = 0$.

Notations

1. The term $(x-a)^n\epsilon(x)$ where $\epsilon(x) \rightarrow 0$ as $x \rightarrow a$ is often denoted as $o_a((x-a)^n)$, this means that the function $x \mapsto o_a((x-a)^n)$ satisfies the property:

$$\lim_{x \rightarrow a} \frac{o_a((x-a)^n)}{(x-a)^n} = 0.$$

2. The remainder $x^n\epsilon(x)$ of limited development of a function f at the point 0 of order n can also be written as $x^n\epsilon(x) = o(x^n)$.

Proposition 2. If a function f has a limited development at a point, then this limited development is unique.

Proposition 3. - If f is an even function, then the polynomial part of its limited development at 0 contains only monomials of even degrees (i.e., $x^0, x^2, x^4, \dots, x^{2n}, n \in \mathbb{N}$).

- If f is an odd function, then the polynomial part of its limited development at 0 contains only monomials of odd degrees (i.e., $x, x^3, x^5, \dots, x^{2n+1}, n \in \mathbb{N}$).

Example 3. On considère la fonction $f(x) = \cos(x)$, qui est de classe $C^\infty(\mathbb{R})$. It admits a limited development at 0 of order 5.

$$f(x) = \cos(x), \quad f'(x) = -\sin(x), \quad f''(x) = -\cos(x), \quad f'''(x) = \sin(x), \quad f^{(4)}(x) = \cos(x), \quad f^{(5)}(x) = -\sin(x).$$

Evaluating each at $x = 0$, we obtain: $f(0) = 1, \quad f'(0) = 0, \quad f''(0) = -1, \quad f'''(0) = 0, \quad f^{(4)}(0) = 1, \quad f^{(5)}(0) = 0$.

Thus, the limited development of $\cos(x)$ at $x = 0$ to order 5 is:

$$f(x) = f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \frac{f'''(0)}{3!}x^3 + \frac{f^{(4)}(0)}{4!}x^4 + \frac{f^{(5)}(0)}{5!}x^5 + o(x^6)$$

By substituting the calculated values :

$$f(x) = 1 + 0 \cdot x + \frac{-1}{2!}x^2 + 0 \cdot x^3 + \frac{1}{4!}x^4 + 0 \cdot x^5 + o(x^6) = 1 - \frac{x^2}{2} + \frac{x^4}{24} + o(x^6)$$

Since f is an even function, the polynomial part of its Taylor expansion only contains terms of even degrees, as stated in the proposition.

1.3 The Limited development of Common functions at 0 up to Order n

- $e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots + \frac{x^n}{n!} + o(x^n)$
- $\ln(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} + \dots + (-1)^{n+1} \frac{x^n}{n} + o(x^n)$
- $\forall \alpha \in \mathbb{R}, \quad (1+x)^\alpha = 1 + \alpha x + \frac{\alpha(\alpha-1)}{2!}x^2 + \dots + \frac{\alpha(\alpha-1)\dots(\alpha-n+1)}{n!}x^n + o(x^n)$
- $\frac{1}{1+x} = 1 - x + x^2 - x^3 + \dots + (-1)^n x^n + o(x^n)$
- $\frac{1}{1-x} = 1 + x + x^2 + x^3 + \dots + x^n + o(x^n)$
- $\sin(x) = x - \frac{x^3}{3!} + \frac{x^5}{5!} + \dots + (-1)^n \frac{x^{2n+1}}{(2n+1)!} + o(x^{2n+2})$
- $\cos(x) = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} + \dots + (-1)^n \frac{x^{2n}}{(2n)!} + o(x^{2n+1})$
- $\tan(x) = x + \frac{x^3}{3} + \frac{2}{15}x^5 + \frac{17}{315}x^7 + o(x^8)$
- $\cosh(x) = 1 + \frac{x^2}{2!} + \frac{x^4}{4!} + \dots + \frac{x^{2n}}{(2n)!} + o(x^{2n+1})$
- $\sinh(x) = x + \frac{x^3}{3!} + \frac{x^5}{5!} + \dots + \frac{x^{2n+1}}{(2n+1)!} + o(x^{2n+2})$
- $\tanh(x) = x - \frac{x^3}{3} + \frac{2}{15}x^5 - \frac{17}{315}x^7 + o(x^8)$

Proposition 4. *A function f has a limited development near to a point a if and only if the function $f(x + a)$ has a limited development near to a point 0 .*

Example 4. *Calculate the $LD_n(1)$ of the function $f(x) = e^x$.
We pose $h = x - 1$, if x is near to 1 , then h is near to 0 . Then*

$$\begin{aligned} e^x &= e^{x-1+1} \\ &= e^1 \times e^{x-1} \\ &= e \times e^h \\ &= e \left(1 + h + \frac{h^2}{2!} + \dots + \frac{h^n}{n!} + h^n \varepsilon(x) \right) \\ &= e + \frac{e}{1!}(x-1) + \frac{e}{2!}(x-1)^2 + \dots + \frac{e}{n!}(x-1)^n + o(x-1)^n, \end{aligned}$$

2 Operations on limited developments

2.1 Sum and Product

Definition 2. *Let $n, p \in \mathbb{N}^*$ with $n < p$. Truncating a polynomial of degree p to order n means keeping only the monomials of degrees $\leq n$.*

Example 5. *1. Truncate the polynomial $P(x) = x^8 + 2x^7 - 3x^5 + 2x^4 - x^3 + x^2 + 1$ to order 5. We will denote the obtained polynomial by $T_5(x)$.*

$$T_5(x) = 1 + x^2 - x^3 + 2x^4 - 3x^5$$

2. Truncate the polynomial $(3x+4x^2)(1+x+x^3)$ to order 2. We will denote the obtained polynomial by $T_2(x)$. We have:

$$(3x + 4x^2)(1 + x + x^3) = 3x(1 + x + x^3) + 4x^2(1 + x + x^3) = 3x + 7x^2 + 4x^3 + 3x^4 + 4x^5$$

Then

$$T_2(x) = 3x + 7x^2$$

Proposition 5. *Consider two functions f and g that admit limited development at 0 of order n :*

$$f(x) = c_0 + c_1x + \dots + c_nx^n + x^n\varepsilon_1(x) \quad \text{and} \quad g(x) = d_0 + d_1x + \dots + d_nx^n + x^n\varepsilon_2(x),$$

with $\lim_{x \rightarrow 0} \varepsilon_1(x) = \lim_{x \rightarrow 0} \varepsilon_2(x) = 0$.

Then:

- *$f + g$ admits a limited development at 0 of order n , given by:*

$$f(x) + g(x) = (c_0 + d_0) + (c_1 + d_1)x + \cdots + (c_n + d_n)x^n + x^n \varepsilon(x),$$

where $\lim_{x \rightarrow 0} \varepsilon(x) = 0$.

- $f \times g$ admits a limited development at 0 of order n , given by:

$$f(x) \times g(x) = T_n(x) + x^n \varepsilon(x),$$

where $\lim_{x \rightarrow 0} \varepsilon(x) = 0$, and $T_n(x)$ is the polynomial

$$(c_0 + c_1x + \cdots + c_nx^n) \times (d_0 + d_1x + \cdots + d_nx^n)$$

truncated to order n .

Example 6. Calculate the limited development at 0 of order 2 of the function $f(x) = e^x + \ln(1+x)$. The functions e^x and $\ln 1+x$ are C^∞ near 0, so we can write their limited development at 0 of order 2:

$$e^x = 1 + x + \frac{x^2}{2!} + o(x^2)$$

and

$$\ln(1+x) = x - \frac{x^2}{2} + o(x^2).$$

then

$$f(x) = \left(1 + x + \frac{x^2}{2!} + o(x^2)\right) + \left(x - \frac{x^2}{2} + o(x^2)\right) = 1 + 2x + o(x^2)$$

Example 7. Calculate the limited development at 0 of order 2 of the function $f(x) = \cos(x) \times \sqrt{1+x}$.

The functions $\cos(x)$ and $\sqrt{1+x}$ are C^∞ near 0, so we can write their limited development at 0 up to order 2:

$$\cos(x) = 1 - \frac{x^2}{2!} + o(x^2) = 1 - \frac{x^2}{2} + o(x^2).$$

and

$$\sqrt{1+x} = (1+x)^{\frac{1}{2}} = 1 + \frac{1}{2}x + \frac{\frac{1}{2}(\frac{1}{2}-1)}{2!}x^2 + o(x^2)$$

then

$$\left(1 - \frac{x^2}{2} + o(x^2)\right) \times \left(1 + \frac{1}{2}x - \frac{1}{8}x^2 + o(x^2)\right) = 1 + \frac{1}{2}x - \frac{5}{8}x^2 + o(x^2)$$

By truncating the result at order 2, we obtain

$$f(x) = 1 + \frac{1}{2}x - \frac{1}{8}x^2 + o(x^2)$$

2.2 Composition

Proposition 6. *We consider two functions f and g that admit a limited development at 0 up to order n :*

$$f(x) = c_0 + c_1x + \cdots + c_nx^n = C(x) + x^n\varepsilon_1(x) \quad \text{and} \quad g(x) = d_0 + d_1x + \cdots + d_nx^n = D(x) + x^n\varepsilon_2(x),$$

with $\lim_{x \rightarrow 0} \varepsilon_1(x) = \lim_{x \rightarrow 0} \varepsilon_2(x) = 0$.

If g admits a limited development at 0 up to order n and f admits a at $g(0)$ up to order n , then $f \circ g$ admits a limited development at 0 up to order n . This expansion is obtained by substituting the limited development of g into that of f and retaining only terms of degree $\leq n$.

Example 8. *Let the functions $f(x) = e^x$ and $g(x) = \sin(x)$. Calculate the $LD_2(0)$ of the composition $f(g(x)) = e^{\sin(x)}$.*

The $LD_2(0)$ of $f(x)$ and $g(x)$ are

$$f(x) = e^x = 1 + x + \frac{x^2}{2} + o(x^2)$$

$$g(x) = x - \frac{x^3}{6} + o(x^3).$$

Then

$$f(g(x)) = e^{\sin(x)} = 1 + \left(x - \frac{x^3}{6} + o(x^3)\right) + \frac{1}{2} \left(x - \frac{x^3}{6} + o(x^3)\right)^2 + o\left(\left(x - \frac{x^3}{6} + o(x^3)\right)^2\right)$$

By truncating the result at order 2, we obtain

$$f(g(x)) = 1 + x - \frac{x^3}{6} + \frac{1}{2} \left(x^2 - \frac{x^4}{3} + o(x^4)\right) + o(x^2) = 1 + x + \frac{x^2}{2} + o(x^2)$$

2.3 Quotient

Consider two functions f and g that admit a limited development at 0 up to order n :

$$f(x) = c_0 + c_1x + \cdots + c_nx^n + x^n\varepsilon_1(x) \quad \text{and} \quad g(x) = d_0 + d_1x + \cdots + d_nx^n + x^n\varepsilon_2(x),$$

with $\lim_{x \rightarrow 0} \varepsilon_1(x) = \lim_{x \rightarrow 0} \varepsilon_2(x) = 0$.

To calculate the limited development of the quotient $\frac{f}{g}$, we will use the limited development of

$$\frac{1}{1+u} = 1 - u + u^2 - u^3 + \cdots + (-1)^n u^n + o(u^n),$$

We have three possible cases:

Case 1: If $d_0 = 1$, we set $u = d_1x + \cdots + d_nx^n + x^n\varepsilon_2(x)$, and the quotient can be written as

$$\frac{f}{g} = f \times \frac{1}{1+u}.$$

Case 2: If $d_0 \neq 0$ and $d_0 \neq 1$, then we reduce to the previous case by rewriting

$$\frac{1}{g(x)} = \frac{1}{d_0} \times \frac{1}{1 + \frac{d_1}{d_0}x + \dots + \frac{d_n}{d_0}x^n + x^n \frac{\varepsilon_2(x)}{d_0}}.$$

Case 3: If $d_0 = 0$, then we factor by x^k (for some k) to reduce to one of the previous cases.

Example 9. We aim to find the $LD_4(0)$ of the function $f(x) = \frac{x}{\sin(x)}$.
The limited development of $\sin(x)$ at $x = 0$ is given by:

$$\sin(x) = x - \frac{x^3}{6} + \frac{x^5}{120} + o(x^5)$$

then

$$f(x) = \frac{x}{\sin(x)} = \frac{x}{x - \frac{x^3}{6} + \frac{x^5}{120} + o(x^5)} = \frac{1}{1 - \frac{x^2}{6} + \frac{x^4}{120} + o(x^4)}$$

Using the expansion for $\frac{1}{1+u}$

$$\frac{1}{1+u} = 1 - u + u^2 + o(x^2)$$

where

$$u = -\frac{x^2}{6} + \frac{x^4}{120} + o(x^4).$$

We find

$$f(x) = 1 - \left(-\frac{x^2}{6} + \frac{x^4}{120}\right) + \left(-\frac{x^2}{6} + \frac{x^4}{120}\right)^2 + o(x^4)$$

By truncating the result at order 4, we obtain

$$f(x) = 1 - \left(-\frac{x^2}{6} + \frac{x^4}{120}\right) + \left(-\frac{x^2}{6}\right)^2 + o(x^4)$$

$$f(x) = 1 + \frac{x^2}{6} + \frac{11x^4}{120} + o(x^4)$$

2.4 Integration

Let F be a primitive of f . The function F has a limited development at a up to order $n+1$, which is written as:

$$F(x) = F(a) + c_0(x-a) + c_1 \frac{(x-a)^2}{2!} + \dots + c_n \frac{(x-a)^{n+1}}{(n+1)!} + (x-a)^{n+1} \theta(x)$$

where $\lim_{x \rightarrow a} \theta(x) = 0$.

This means that we integrate the polynomial part term by term to obtain the Taylor expansion of $F(x)$, starting from the constant $F(a)$.

Example 10. Let $f(x) = \arctan(x)$ be defined on \mathbb{R} . Let's find its limited development at 0. We have $f'(x) = \frac{1}{1+x^2}$, the $LD_n(0)$ of $\frac{1}{1+x^2}$ at 0 is:

$$f'(x) = \frac{1}{1+x^2} = 1 - x^2 + x^4 - \dots + (-1)^n x^{2n} + o(x^{2n})$$

Now, integrate $f'(x)$ to find $f(x)$:

$$f(x) = \int (1 - x^2 + x^4 - \dots + (-1)^n x^{2n} + o(x^{2n})) dx$$

we get:

$$f(x) = x - \frac{x^3}{3} + \frac{x^5}{5} - \dots + (-1)^n \frac{x^{2n+1}}{2n+1} + o(x^{2n+1})$$

3 Applications of Limit Developments

3.1 Limit Calculation

We seek to calculate $\lim_{x \rightarrow a} f(x)$. If f admits a limited development at a to the order n , then

$$f(x) = c_0 + c_1(x-a) + \dots + c_n(x-a)^n + (x-a)^n \varepsilon(x)$$

and therefore,

$$\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} (c_0 + c_1(x-a) + \dots + c_n(x-a)^n + (x-a)^n \varepsilon(x)).$$

Example 11. Calculate $\lim_{x \rightarrow 0} \frac{\sin(x) - x}{x(\cos(x) - 1)}$.

We can see that this limit is an indeterminate form. We know the $DL_3(0)$ of $\sin(x)$ and $\cos(x)$ in 0

$$\begin{aligned} \sin(x) - x &= \frac{x^3}{3!} + x^3 \varepsilon(x), \\ x(\cos(x) - 1) &= \frac{x^3}{2!} + x^3 \varepsilon(x), \end{aligned}$$

by substituting we have

$$\lim_{x \rightarrow 0} \frac{\sin(x) - x}{x(\cos(x) - 1)} = \lim_{x \rightarrow 0} \frac{\frac{x^3}{3!}}{\frac{x^3}{2!}} = \lim_{x \rightarrow 0} \frac{\frac{1}{6}}{\frac{1}{2}} = \frac{1}{3}.$$

3.2 Continuity and Differentiability from a limited development

Let $f(x)$ be a function defined on an interval I , except at $x = 0$. If f admits a first-order limited development at $x = 0$, meaning $f(x) = a_0 + a_1x + x\epsilon(x)$; then the following results hold:

1. **Extension by Continuity:** we can extend the function to be continuous at $x = 0$ by defining: $f(0) = a_0$

2. **Differentiability at $x = 0$:** This extension guarantees that $f(x)$ is differentiable at $x = 0$, with $f'(0) = a_1$

3. **Equation of the Tangent Line:** The equation of the tangent line to $f(x)$ at $x = 0$ is: $y = a_0 + a_1x$

Example 12. Let $f(x) = \frac{e^x - 1}{x}$ with $e^x = 1 + x + \frac{x^2}{2!} + o(x^2)$, then

$$f(x) = \frac{x + \frac{x^2}{2} + x^2\epsilon(x)}{x} = 1 + \frac{x}{2} + x^2\epsilon(x)$$

1. We can extend f by continuity at $x = 0$ by posing $f(0) = 1$.
2. The function is differentiable at $x = 0$ with $f'(0) = \frac{1}{2}$
3. The equation of the tangent line to $f(x)$ at $x = 0$ is $y = 1 + \frac{x}{2}$.